

AD 689045

U.S. ARMY FOREIGN SCIENCE AND TECHNOLOGY CENTER



A METHOD OF OBTAINING LUMINESCENT SERA

COUNTRY: USSR

TECHNICAL TRANSLATION

Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.

Reproduction of this document in whole or in part by non Department of Defense agencies

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

TECHNICAL TRANSLATION

FSTC-HT-23-754-68

A Method of Obtaining Luminescent Sera

by

Yu.Ye. Sklyar, G.I. Mikhaylov,
K.L. Shakhanina & V.A. Chibisova

All-Union Scientific Research Institute of
Chemical Reagents and Extra Pure Chemical Substances

and

Institute of Epidemiology and Microbiology named
for N. F. Gamaley

Source: IZOBRETENIYA, PROMYSHLENNYYE OBRAZTSY, TOVARNYYE
ZNAKI

(Inventions, Industrial Samples, Trademarks)

No. 8, pp 69-70, 1967

USSR

Translated for FSTC by ACSI

This translation is an unedited rendition of the original foreign text. Statements or theories advocated or implied are those of the source and do not reflect the position or opinion of the US Army Foreign Science and Technology Center. This translation is published with a minimum of copy editing and graphics preparation in order to expedite the dissemination of information. Requests for additional copies of this document should be addressed to the Defense Documentation Center, Cameron Station, Alexandria, Virginia, ATTN: OSR-2.

A METHOD OF OBTAINING LUMINESCENT SERA

1

A method is known of obtaining luminescent sera by means of attaching to them fluorescent isocyanate, dimethylaminonaphthalenesulfochloride and other fluorochromes.

The proposed method consists of the use of dichlorotriazinylaminofluorescein as the label, which allows one to replace the difficultly available dye.

Example 1. 1.77g of cyanurichloride is dissolved in 10ml of anhydrous acetone, cooled to 0°C. and with energetic stirring one adds dropwise a solution of 2.91g of aminofluorescein I in 100ml of anhydrous acetone. This is mixed at 0°C for two hours, and having precipitated a light yellow precipitate is filtered and washed with 10ml of anhydrous acetone.

Yield 3.77g (84.3%). m.p. 300°C.

Example 2. 321.3mg of cyanurichloride in 10ml of anhydrous acetone is cooled to 0°C and with careful stirring one adds dropwise a solution of 514.4mg of aminofluorescein in 50ml of anhydrous acetone. This is mixed for two hours at 0°C, and having precipitated a light yellow precipitate of ohlohydrate of dichlorotriazinylaminofluorescein is filtered and washed with anhydrous acetone.

Yield 622.3mg (79%). m.p. 300°C.

Example 3. 100ml of native serum is precipitated with ammonium sulfate at 40-50% saturation. After six hours the precipitate is isolated on a Buchner funnel, dissolved in distilled water so that the content of protein was 3-4%, and salt is removed by successive passage of the solution through cation-exchange and anion-exchange resins according to the published method.

Dichlorotriazinylaminofluorescein in a quantity of 15mg per lg of protein is pulverized in a agate mortar with a small amount of 0.1M phosphate buffer (pH 8.6) and is added to a 2-2.5% solution of protein in the same buffer. Conjugation occurs after mixing for six hours at room temperature. Free dye is removed on Sephadex G-25 (course), 0.01M phosphate buffer (pH 7.5). Protein with a content of fluorescein greater than 3.5 moles per mole of protein, causing nonspecific luminescent background, are separated by adsorption on DEAE-cellulose at pH 7.5 (2g DEAE-cellulose with moisture of 65% per 10ml of conjugate). To the serum one adds merthiolate (1:10,000) and sodium chloride to the physiological concentration. After three days the serum is centrifuged (40,000 rpm) and is subjected to lyophilic drying.

Object of the Invention

The method of obtaining luminescent sera by means of labeling of immune antibodies by a fluorochrome, is distinguished by the purpose of replacing the deficient fluorescein-thiocyanate, and using dichlorotriazinylaminofluorescein as the label.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body, abstract and indexing annotation must be entered when the overall report is classified)

| | | | |
|---|--|---|------------------------|
| 1. ORIGINATING ACTIVITY (Corporate author) Foreign Science and Technology Center US Army Materiel Command Department of the Army | | 2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED | |
| | | 2b. GROUP | |
| 3. REPORT TITLE A Method of Obtaining Luminescent Sera | | | |
| 4. DESCRIPTIVE NOTES (Type of report and inclusive dates) - Translation | | | |
| 5. AUTHOR(S) (First name, middle initial, last name) Yu. Ye. Sklyar, G. I. Mikhaylov, K.L. Shakhanina & V.A. Chibisova | | | |
| 6. REPORT DATE 5 MAY 1969 | | 7a. TOTAL NO. OF PAGES 2 | 7b. NO. OF REFS N/A |
| 8a. CONTRACT OR GRANT NO. A. PROJECT NO. c. 9223628 2301 d. SMUFD-PR-I | | 8b. ORIGINATOR'S REPORT NUMBER(S) FSTC-HT-23-754-68 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) ACSI Control Number J 6316 | |
| 10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited. | | | |
| 11. SUPPLEMENTARY NOTES | | 12. SPONSORING MILITARY ACTIVITY US Army Foreign Science and Technology Center | |
| 13. ABSTRACT The method of obtaining luminescent sera by means of labeling of immune antibodies by a fluorochrome, is distinguished by the purpose of replacing the deficient fluorescein thiocyanate, and using dichlorotriazinylaminofluorescein as the label. | | | |

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

UNCLASSIFIED

Security Classification

| 14. | KEY WORDS | LINK A | | LINK B | | LINK C | |
|-----|--|--------|----|--------|----|--------|----|
| | | ROLE | WT | ROLE | WT | ROLE | WT |
| | Luminescent sera Fluorescent isocyanate Dimethylaminonaphthalenesulfochloride Fluorochromes | | | | | | |